

FOR ANIMAL USE ONLY

ProndiVAX

PULPY KIDNEY

Reg. No. G2548 (Act 36/1947)

INDICATIONS:

For the active immunisation of Sheep and Goats against **Pulpy Kidney disease** (enterotoxaemia).

STORAGE INSTRUCTIONS:

- STORE BETWEEN 2 °C AND 8 °C.
- DO NOT FREEZE.
- PROTECT FROM LIGHT.

COMPOSITION:

Off-white suspension containing formalized, alum-precipitated *Clostridium perfringens* type D (epsilon) toxoids.

WARNINGS:

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF VACCINATION.
- Usually no marked reaction follows vaccination although a transient swelling may appear at the site of inoculation, and some animals may show a moderate rise in temperature for one or two days.
- Keep out of reach of children, uninformed persons and animals.
- Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED:

- Shake bottle well before use and repeat from time to time during the vaccination process.
- Part-used containers should be disposed of at the end of each day's operation since under field conditions it may be difficult to avoid accidental contamination of the vaccine.
- If possible use a separate, sharp, sterile needle for each animal.
- Administer the vaccine subcutaneously on the side of the neck midway between the shoulder and the head.
- Goats should be vaccinated under the loose skin underneath the tail.
- Do not vaccinate Goats in the neck.

- Sheep and Goats not previously vaccinated should be given two injections of vaccine, with an interval of 4 - 6 weeks.
- Lambs and kids can be immunized from 3 months of age. Administer a second dose 4 - 6 weeks later.
- An annual booster injection should be given to the whole flock 2 - 6 weeks before the ewes are due to lamb.
- Lambs born from ewes vaccinated before lambing are protected for 12 - 16 weeks by the antibodies passed in the colostrum.
- It is sound practice to reinoculate vaccinated Sheep shortly before the period of greatest risk.

Note: Lambs and kids younger than 2 months and animals in poor condition, infested with parasites or which are suffering from other diseases will not immunise well.

DOSE:

Sheep and Goats of all ages
1 ml subcutaneously.

PRESENTATION:

100 ml (100 doses) multi-dose vials.
250 ml (250 doses) multi-dose vials.

Registration holder:

Animal Health Registration World Wide (Pty) Ltd
Co. Reg. No. 1998/17802/07

Manufactured by:

Prondil SA, Montevideo Uruguay
www.prondil.com

Distributed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009, Faerie Glen, 0043 R.S.A.
Tel: +27 (0)12 817 9060
Web: www.afrivet.co.za
Helpline: 0860 VEEARTS (0860 833 2787)

ProndiVAX

PULPY KIDNEY

Reg. nr. G2548 (Wet 36/1947)

INDIKASIES:

Vir die aktiewe immunisering van Skape en Bokke teen **Bloednier**.

BERGINGSaanwysings:

- BERG TUSSEN 2 °C EN 8 °C.
- MOET NIE BEVRIES NIE.
- BESKERM TEEN LIG.

SAMESTELLING:

Naaswit suspensie wat geformaliseerde, aluin-gepresipiteerde *Clostridium perfringens* tipe D (epsilon) toksioïedes bevat.

WAARSKUWINGS:

- MOET NIE DIERE BINNE 7 DAE NA INENTING VIR MENSLIKE VERBRUIK SLAG NIE.
- Geen opmerkbare reaksies word gewoonlik na enting opgemerk maar swelling mag by die plek van toediening voorkom en party diere mag 'n matige styging in temperatuur vir een of twee dae toon.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie entstof breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEbruIKSAANWYSINGS - GEbruIK SLEGS SOOS AANGEDUI:

- Skud die bottel goed voor gebruik en herhaal van tyd tot tyd gedurende die inenting proses.
- Houers wat gedeeltelik gebruik is, moet na daaglikse gebruik vernietig word aangesien dit onder veld toestande moeilik is om te verhoed dat die entstof besmet word.
- Indien moontlik gebruik 'n aparte, skerp, steriele naalde vir elke dier.
- Onderhuidse toediening aan die kant van die nek halfpad tussen die kop en skouer.
- Bokke moet in die los vel onder die stert ingespuut word.
- Moenie Bokke in die nek spuit nie.
- Skape en Bokke wat nooit

voorheen geënt was nie benodig twee inspuitings entstof, met 'n tussenpose van 4 - 6 weke.

- Lammers kan van 3 maande ouderdom geïmmuniseer word. 'n Tweede inenting moet 4 - 6 weke later gegee word.
- 'n Jaarlikse skraagdosis moet 2 - 6 weke voor lam aan die hele kudde toegedien word.
- Ooie wat voor lam geënt word sal immuniteit oordra deur middel van biesmelk en lammers sal beskerm wees vir 12 - 16 weke.
- Dit is raadsaam om Skape, kort voor die grootste risiko periode, te her-ent.

Nota: Lammers jonger as 2 maande en diere in swak kondisie, met parasiete besmet of wat aan ander siektes lei sal nie goeie immunisering opbou nie.

DOSIS:

Skape en Bokke van alle ouderdomme: 1 ml onderhuids.

AANBIEDING:

100 ml (100 dosisse) multi-dosis bottels.

250 ml (250 dosisse) multi-dosis bottels.

Registrasiehouer:

Animal Health Registration World Wide (Edms) Bpk Mpy. Reg. nr. 1998/17802/07

Vervaardig deur:

Prondil SA, Montevideo Uruguay
www.prondil.com

Versprei deur:

Afrivet Business Management (Edms) Bpk Mpy. Reg. nr. 2000/011263/07
Posbus 2009, Faerie Glen, 0043 R.S.A.
Tel: +27 (0)12 817 9060
Web: www.afrivet.co.za
Hulplyn: 0860 VEEARTS (0860 833 2787)